



Product Name: ICON POUR-ON IVERMECTIN ENDECTOCIDE FOR CATTLE
APVMA Approval No: 92264/136301

Label Name:	ICON POUR-ON IVERMECTIN ENDECTOCIDE FOR CATTLE
-------------	--

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	10 mg/mL IVERMECTIN
-------------------------	---------------------

Claims:	<p>For the treatment and control of ivermectin sensitive internal and external parasites of beef and dairy cattle.</p> <p>ICON POUR-ON IVERMECTIN ENDECTOCIDE FOR CATTLE contains ivermectin, a member of the macrocyclic lactones family of chemicals. It is effective against sensitive strains of the following internal parasites:</p> <p>For the treatment and control of adult and some immature stages of:</p> <ul style="list-style-type: none">Gastro-intestinal RoundwormsBarbers Pole Worm (<i>Haemonchus placei</i>)Small Brown Stomach Worm (<i>Ostertagia ostertagi</i>) (including inhibited stages)Small Intestinal Worm (<i>Cooperia oncophora</i>, <i>Cooperia punctata</i>)Stomach Hair Worm (<i>Trichostrongylus axei</i>)Hookworm (<i>Bunostomum phlebotomum</i>)Large Bowel Worm (<i>Oesophagostomum radiatum</i>)Whipworm (<i>Trichuris</i> spp.)Thin Necked Intestinal Worm (<i>Nematodirus spathiger</i>)Intestinal Threadworm (<i>Strongyloides papillosus</i>)Lungworms (<i>Dictyocaulus viviparus</i>) <p>Sucking and Biting Lice</p> <ul style="list-style-type: none">Long Nosed Sucking Lice (<i>Linognathus vituli</i>)Short Nosed Sucking Lice (<i>Haematopinus eurysternus</i>)Biting Lice (<i>Bovicola bovis</i>)Little Sucking Lice (<i>Solenopotes capillatus</i>) <p>Mites (<i>Chorioptes bovis</i>, <i>Sarcoptes scabiei</i> var <i>bovis</i>)</p> <p>Buffalo Fly (<i>Haematobia irritans exigua</i>)</p> <p>Cattle Tick (<i>Rhipicephalus (Boophilus) microplus</i>)</p>
---------	---

Maximum tick efficacy is achieved 4-5 days after treatment. Engorged female ticks that drop from cattle in the initial days after treatment may lay viable eggs. This should be taken into account when planning a strategic tick control program. ICON POUR-ON should be alternated with conventional dips or pour-ons when the program requires repeat treatments.

Sustained Activity

At the recommended dose rate, ICON POUR-ON effectively controls infection with *Ostertagia* spp. (Small brown stomach worm) acquired up to 14 days after treatment, with *Dictyocaulus viviparus* (Lungworm) acquired up to 28 days after treatment and *Haematobia irritans exigua* (Buffalo Fly) up to 21 days after treatment.

Resistance may develop to any chemical.

Net Contents: 250 mL, 1 L, 2.5 L, 3 L, 5 L, 5.5 L, 15 L, 20 L

Directions for Use:

Restrains: Re-treatment interval: DO NOT re-treat animals for 21 days after last treatment.

Contraindications:

Precautions: Rainfall on cattle less than 2 hours after dosing may result in reduced efficacy.

Side Effects: Temporary hair loss occasionally occurs along the application site. This resolves without treatment.

Dosage and Administration: ICON POUR-ON dose rate is 1 mL/20 kg body weight. Apply along the backline of the animal.

<table>

Cattle heavier than 650 kg should be dosed at 1 mL per 20 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

This section contains file attachment.

General Directions:

Withholding Periods: MEAT: DO NOT USE less than 21 days before slaughter for human consumption.
MILK: Zero (0) days.

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. Before using this product, confirm the current ESI from Alleva Animal Health Ltd on 1800 255 382 or the APVMA website (www.apvma.gov.au/residues).
Safety Directions:	Poisonous if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. When using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
First Aid Warnings:	
Additional User Safety:	Additional information is listed in the safety data sheet.
Environmental Statements:	ENVIRONMENTAL PROTECTION Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.
Disposal:	[250 mL, 1 L] Dispose of container by wrapping with paper and putting in garbage. [Other pack sizes] Triple-rinse container into the medicated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
Storage:	Store below 30°C (room temperature).

Liveweight (kg)	Dose volume (mL)
51-100	5
101-150	7.5
151-200	10
201-250	12.5
251-300	15
301-350	17.5
351-400	20
401-450	22.5
451-500	25
501-550	27.5
551-600	30
601-650	32.5